Spotted Lanternfly



Quick Take What:

The spotted lanternfly—a destructive, invasive plant hopper attacks many hosts including grapes, apples, stone fruits, walnut, willow, and tree of heaven—has been confirmed in New Castle County. Delaware is the second state to have found the insect, which was first detected in Pennsylvania in 2014.

Significance:

Agriculture is Delaware's #1 industry, providing 75,000 jobs and \$12.7 billion of economic activity, including \$1.2 billion in direct sales from our 2,158 family farms.

This insect is a potential threat to several important commercial crops including grapes, peaches, apples, and timber. Early detection is vital for the protection of Delaware businesses and agriculture.



What is Spotted Lanternfly?

The spotted lanternfly, *Lycorma delicatula*, is a plant hopper native to China, India, and Vietnam, that belongs to the order Homoptera, family Fulgoridae.

The spotted lanternfly adult is 1" long and 1/2" wide at rest. The forewings are gray with black spots, and the hind wings are red with black spots. The head and legs are black, and the abdomen is yellow with broad black bands. Immature stages are small, round, and black with white spots, and develop red patches as they grow.

Officials believe the spotted lanternfly must feed on the tree of heaven (*Ailanthus altissima*) to reproduce. However, the insect can also be found feeding on other plants and trees. Egg masses will be laid on the trunk, branches, and limbs of medium to large trees, often in the upper reaches of the canopy. In springtime, nymphs will hatch and move off the trees and search for new hosts, including several agricultural crops, such as grapes, apples, and peaches.

#HitchHikerBug

The spotted lanternfly is a distinct insect in both the nymph and adult phases. The egg masses can be hard to spot making it easy to transport them on vehicles and other outdoor items.

Based on the proximity to Pennsylvania and the ease with which the spotted lanternfly can "hitch-hike," officials believe this could be how the insect was introduced to Delaware. Therefore, Delawareans and visitors must inspect their vehicles and anything they may be transporting, including trailers, stone, metal, or anything that has spent time outdoors, and remove and destroy any attached egg mass. They have been seen hitching rides in and on cars and clothing. It is important to check for any sign of the spotted lanternfly before traveling to and from surrounding states, especially Pennsylvania, New Jersey, Maryland, and Virginia.






What to do...

If you find a spotted lanternfly, please capture and destroy this invasive pest.

Positively identified spotted lanternfly should be destroyed whenever possible with registered insecticides, and discarded. Egg masses can scraped into a zipped plastic bag containing rubbing alcohol or an alcohol-based hand sanitizer, then crush the eggs and discard.



SPOTTING GUIDE

SPOTTED LANTERNFLY LIFE STAGES









From April to November, spotted lanternfly grow and change rapidly.

Throughout the spring and summer, be on the lookout for the different life stages of spotted lanternfly.

It's the Law!

The Delaware Department of Agriculture has the authority under the Plant Pest Law (Title 3, Chapter 11 § 1103) to issue orders for any control measures it deems necessary to prevent the spread of dangerously injurious plant pests. Upon knowledge of the existence of such plant pest, such as spotted lanternfly, the Department shall notify the person(s), owner(s) or the tenant(s) in possession of the premises in question of the existence of this pest and to implement the prescribed control measures within the prescribed time limit, otherwise the individual will be subject to civil penalties.

If the person associated with or owners of the plants infested with spotted lanternfly are found to neglect, fail or refuse to apply the control measures prescribed by the Department, in the manner or at the times ordered and directed by the Department, then the Department may cause the prescribed control measures to be applied at the expense of that person.

What I Am Not...









Boxelder Bug Leopard Moth

Tiger Moth

Delaware SLF Quarantine

In 2019, Delaware established a spotted lanternfly quarantine due to detecting established populations in New Castle County, Delaware. A quarantine is a regulation to slow the spread of the spotted lanternfly, and means that any material or object that could harbor the pest cannot be moved without taking precautions to prevent the spread. The general public must complete a residential compliance checklist indicating that regulated articles have been inspected and found free of any life stages of spotted lanternfly. Any business requiring the movement of any regulated item within a spotted lanternfly quarantine area must have a permit.

In October 2020, the Delaware Department of Agriculture expanded the quarantine to include Kent County, and subsequently quarantined Sussex County in July 2022 when established populations were found. For the most up-to-date details and information regarding the Delaware Spotted Lanternfly Quarantine, and to apply for a permit, visit https://de.gov/hitchhikerbug.

Signs and Symptoms of SLF

Spotted lanternfly feeds on sap from trees using their piercing-sucking mouthparts, causing noticeable weeping wounds on susceptible trees, leaving a grayish or black sticky trail along the trunk. This sap will attract other insects to feed, notably wasps and ants.

Spotted lanternfly also secrete a high-sugar substance called honeydew, which can fall and coat leaves and objects below feeding locations. Not only is it annoying for homeowners, the honeydew causes mold growth on plants that prevents photosynthesis and weakens the plant. It also attracts stinging insects, looking to feed on the honeydew,

Branch dieback, wilting and plant death is a common symptom of heavy spotted lanternfly feedings.

So You Have Tree of Heaven?

If you have found tree of heaven on your property, it is time to make a plan!

- Identify which trees are female.
 These trees will have the winged seedpods. Female trees should be removed from your property to eliminate them as a potential food source for spotted lanternfly.
- 2. Create "trap trees."

Trap trees are male tree of heaven that do not have seeds. Leave 10% of the largest male trees standing and treat them with a registered systemic insecticide to target spotted lanternflies that land and feed on them. This chemical should be applied yearly in the springtime.

3. Time to cut and treat

Now that you have your trap trees
picked out, it's time to remove the
remaining tree of heaven.

Tree of heaven is notorious for growing back with a vengeance if it's only cut down. The roots have been known to reach out 26 feet and will send up offshoots if the parent tree is cut down. In order to ensure that the tree is killed, you will need to treat it with a herbicide.

When using herbicides and insecticides, remember to read the product label and be mindful of all label restrictions!



How Do I Know What to Use to Kill Spotted Lanternfly?

When treating for spotted lanternfly, homeowners should look for products labeled for plant hoppers.

Always make sure to follow label directions!

It's most effective to apply systemic insecticides in the spring and summer before adult populations increase.

Once the adult spotted lanternfly is in the environment, use **contact insecticides**.

The Delaware Department of Agriculture maintains a list of approved spotted lanternfly chemical treatment options online under the Managing Spotted Lanternfly tab at:

https://de.gov/hitchhikerbug

Mention of a product name does not constitute a specific endorsement by the Delaware Department of Agriculture or the University of Delaware.

Tree Identification

Proper identification of tree of heaven is important in helping to eradicate this invasive pest. Although tree of heaven is an invasive plant, there are several native trees that are similar-looking. It's important to distinguish these from tree of heaven so that these look-alikes are not removed.

Tree of Heaven is a very hardy tree species that can be found growing almost anywhere because of its tolerance of poor soils. The species can grow to 82 feet, and is usually found in clusters.

- **Leaves**: Compound leaves with compound leaflets. Leaves and twigs have strong "nutty" odor when crushed or cut.
- **Bark**: Gray bark with vertical lines. As the tree gets older, bark becomes darker.
- Fruit: Females have winged seed pods that hang on the tree through the winter.

Sumac is much smaller when fully grown reaching 32 feet, on average. There are two common sumacs found in Delaware—Staghorn and Smooth.

- Leaves: Compound leaves with serrated leaflets.
- **Bark**: Staghorn has a velvety bark much like the antlers of a deer. Smooth sumac has a brown-gray bark that is smooth and will develop scaly ridges with age.
- Fruit: Clusters of round, finely-haired reddish fruit in an upright cluster.

Black walnut is a tree that can grow upwards of 60-80 feet, but has low tolerance for shade and poor soil.

- Leaves: Compound leaves with smooth leaflet margins.
- **Bark**: Brown on the surface, but darker brown when cut. The bark is ridged and furrowed, with a rough diamond pattern.
- **Fruit**: Matures in the fall, growing between 2-2.5 inches round with a hard outer husk over the nut.

For More Information

Delaware Department of Agriculture—Spotted Lanternfly Program 2320 S. DuPont Hwy, Dover, DE 19901
HitchHikerBug@delaware.gov
de.gov/hitchhikerbug



Tree of Heaven (Ailanthus altissima)



Leaves and winged seed pods of tree of heaven.



Leaves and seed head cluster of staghorn sumac.



Leaves and fruit of a black walnut.